

Emtricitabine

Brand Name: Emtriva



Drug Description

Emtricitabine, also referred to as FTC, is a nucleoside reverse transcriptase inhibitor (NRTI). Emtricitabine is the (-) enantiomer of a thio analogue of cytidine; it differs from other cytidine analogues by a fluorine in the 5 position. [1]

HIV/AIDS-Related Uses

Emtricitabine was approved by the FDA on July 2, 2003 for use in combination with other antiretroviral agents for the treatment of HIV infection in adults age 18 and older. Safety and effectiveness in pediatric patients have not been established. In antiretroviral treatment experienced patients, the use of emtricitabine may be considered for adults with HIV that is expected to be susceptible to emtricitabine as assessed by genotypic or phenotypic testing.[2]

Non-HIV/AIDS-Related Uses

Emtricitabine is being studied for the treatment of chronic hepatitis B virus (HBV) infection.[3]

Pharmacology

Emtricitabine, a synthetic nucleoside analogue of cytosine, undergoes phosphorylation by means of cellular enzymes. The product of phosphorylation, emtricitabine 5'-triphosphate, inhibits viral DNA synthesis by competing with the natural substrate deoxycytidine 5'-triphosphate for incorporation into viral DNA and terminating the DNA chain at the point of incorporation.[4]

Emtricitabine is rapidly and extensively absorbed following oral administration, reaching peak plasma concentrations (C_{max}) at 1 to 2 hours post-dose. In one clinical trial, the mean absolute bioavailability of emtricitabine was 93% following multiple doses of the drug. The mean steady state C_{max} was 1.8 plus or minus 0.7 mcg/ml and the area under the plasma concentration-time curve (AUC) over a 24-hour dosing interval was 10.0 plus or minus 3.1 hr mcg/ml. The mean steady state plasma trough concentration 24 hours after an oral dose was 0.09 mcg/ml.[5]

Emtricitabine is in FDA Pregnancy Category B. Animal studies revealed no increased incidences of fetal variations or malformations in mice and rabbits at 60- and 120-fold higher exposures, respectively, than the human exposure at the recommended daily dose. However, there have been no adequate well-controlled studies in pregnant women. Results of animal studies are not always predictive of human response and emtricitabine should be used during pregnancy only if clearly needed. It is not known whether emtricitabine is distributed into human milk.[6]

Emtricitabine is less than 4% bound to plasma proteins and protein binding is independent of drug concentration over a range of 0.02 to 200 mcg/ml. In vitro studies indicate that emtricitabine does not inhibit CYP450 enzymes. Following administration of ¹⁴C-emtricitabine, the dose was 86% recovered in urine and 14% in feces. Thirteen percent was recovered in urine as metabolites, including 3'-sulfoxide diastereomers and 2'-O-glucuronide; no other metabolites were identifiable. The plasma half-life of emtricitabine is approximately 10 hours. Renal clearance of the drug exceeds estimated creatinine clearance, indicating elimination by glomerular filtration and tubular secretion. In patients with renal impairment, C_{max} and AUC were increased.[7]

HIV isolates with reduced susceptibility to emtricitabine have been recovered from some patients treated with emtricitabine alone or in combination with other antiretroviral agents. Viral isolates from 37.5% of patients with virologic failure had reduced susceptibility to emtricitabine. Genotypic analysis revealed the cause as M184V/I mutations in the HIV reverse transcriptase gene. Cross-resistance has been noted among some nucleoside analogues. Emtricitabine-resistant isolates were cross-resistant to lamivudine and zalcitabine but retained susceptibility to abacavir, didanosine, stavudine, tenofovir, and zidovudine, as well as to the nonnucleoside reverse transcriptase inhibitors delavirdine, efavirenz, and nevirapine.[8]

Adverse Events/Toxicity

The most frequently reported adverse effects of

Emtricitabine



Adverse Events/Toxicity (cont.)

emtricitabine are headache, nausea, skin rash, and skin discoloration on palms and soles.[9]

Lactic acidosis and severe hepatomegaly with steatosis, including fatal cases, have been reported with the use of nucleoside analogues alone or in combination, including emtricitabine. In some patients coinfecting with HIV and hepatitis, exacerbation of hepatitis B has been reported after discontinuing treatment with emtricitabine.[10]

Redistribution of body fat, peripheral wasting, facial wasting, breast enlargement, and cushingoid appearance have been observed in patients receiving antiretroviral therapy.[11]

Drug and Food Interactions

Emtricitabine may be administered with or without food: the AUC was unchanged and C_{max} decreased by 29% when the drug was administered with a 1,000-calorie, high-fat meal.[12]

Emtricitabine has been evaluated in healthy volunteers in combination with tenofovir disoproxil fumarate (tenofovir DF), indinavir, famciclovir, and stavudine. Results showed no interactions except for a small increase in plasma trough concentrations of emtricitabine when it is administered concurrently with tenofovir DF.[13] Because elimination of emtricitabine is probably via both glomerular filtration and active tubular secretion, there may be competition for elimination with other compounds that are also renally eliminated.[14]

Contraindications

Emtricitabine is contraindicated in patients with previously demonstrated hypersensitivity to any of the components of the drug product.[15]

Clinical Trials

For information on clinical trials that involve Emtricitabine, visit the ClinicalTrials.gov web site at <http://www.clinicaltrials.gov>. In the Search box, enter: Emtricitabine AND HIV Infections.

Dosing Information

Mode of Delivery: Oral.[16]

Dosage Form: Capsules containing 200 mg of emtricitabine.[17]

The recommended dose of emtricitabine for adults 18 years of age and older is 200 mg once daily. The dosing interval of emtricitabine should be adjusted in patients with baseline creatinine clearance less than 50 ml/min. The dosing interval recommendations are: creatinine clearance 30 to 49 ml/min, 200 mg every 48 hours; creatinine clearance 15 to 29 ml/min, 200 mg every 72 hours; and creatinine clearance less than 15 ml/min, 200 mg every 96 hours.[18]

Storage: Store at 25 C (77 F); excursions permitted to 15 C to 30 C (59 C to 86 F).[19]

Chemistry

CAS Name: (2R-cis)-4-Amino-5-fluoro-1-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]-2(1H)-pyrimidinone[20]

CAS Number: 143491-57-0[21]

Molecular formula: C₈H₁₀F-N₃O₃-S[22]

C 38.86%, H 4.08%, F 7.68%, N 17.00%, O 19.41%, S 12.97%[23]

Molecular weight: 247.25[24]

Melting point: 136 to 140 C[25]

Physical Description: White to off-white powder.[26]

Solubility: Approximately 112 mg/ml in water at 25 C (77 F).[27]

Other Names

2'-Deoxy-5-fluoro-3'-thiacytidine[28]

2-FTC[29]

524W91[30]

Emtricitabine



Other Names (cont.)

BW524W91[31]

dOTFC[32]

FTC[33]

Coviracil[34]

BW 1592[35]

Emtricitabina[36]

5-fluoro-1-[(2R,5S)-2-(hydroxymethyl)-1,3-oxathiolan-5-yl]cytosine[37]

Further Reading

Gulick RM. New antiretroviral drugs. Clin Microbiol Infect. 2003 Mar;9(3):186-93. Review.

Szzech GM, Wang LH, Walsh JP, Rousseau FS. Reproductive toxicology profile of emtricitabine in mice and rabbits. Reprod Toxicol. 2003 Jan-Feb;17(1):95-108.

Richman DD. Antiretroviral activity of emtricitabine, a potent nucleoside reverse transcriptase inhibitor. Antivir Ther. 2001 Jun;6(2):83-8. Review. No abstract available.

Molina JM, Ferchal F, Rancinan C, Raffi F, Rozenbaum W, Sereni D, Morlat P, Journot V, Decazes JM, Chene G. Once-daily combination therapy with emtricitabine, didanosine, and efavirenz in human immunodeficiency virus-infected patients. J Infect Dis. 2000 Aug;182(2):599-602. Epub 2000 Jul 13.

Darque A, Valette G, Rousseau F, Wang LH, Sommadossi JP, Zhou XJ. Quantitation of intracellular triphosphate of emtricitabine in peripheral blood mononuclear cells from human immunodeficiency virus-infected patients. Antimicrob Agents Chemother. 1999 Sep;43(9):2245-50.

Ena J, Pasquau F. Once-a-day highly active antiretroviral therapy: a systematic review. Clin Infect Dis. 2003 May 1;36(9):1186-90. Epub 2003

Apr 21.

Manufacturer Information

Emtricitabine
Gilead Sciences Inc
333 Lakeside Dr
Foster City, CA 94404
(800) 445-3235

Emtriva
Gilead Sciences Inc
333 Lakeside Dr
Foster City, CA 94404
(800) 445-3235

For More Information

Contact your doctor or an AIDSinfo Health Information Specialist:

- Via Phone: 1-800-448-0440 Monday - Friday, 12:00 p.m. (Noon) - 5:00 p.m. ET
- Via Live Help: http://aidsinfo.nih.gov/live_help Monday - Friday, 12:00 p.m. (Noon) - 4:00 p.m. ET

Emtricitabine



References

1. Merck Index - 2001; p. 630
2. Gilead Sciences - Emtriva Prescribing Information, p.6. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
3. Gilead Sciences - Emtriva Status. Available at: <http://www.gilead.com/wt/sec/emtriva>. Accessed 11/26/03.
4. Gilead Sciences - Emtriva Prescribing Information, p.1. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
5. Gilead Sciences - Emtriva Prescribing Information, p.3. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
6. Gilead Sciences - Emtriva Prescribing Information, p.10. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
7. Gilead Sciences - Emtriva Prescribing Information, p.3-4. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
8. Gilead Sciences - Emtriva Prescribing Information, p.2. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
9. Gilead Sciences - Emtriva Prescribing Information, p.10-11. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
10. Gilead Sciences - Emtriva Prescribing Information, p.8. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
11. Gilead Sciences - Emtriva Prescribing Information, p.9. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
12. Gilead Sciences - Emtriva Prescribing Information, p.3. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
13. Gilead Sciences - Emtriva Prescribing Information, p.5. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
14. Gilead Sciences - Emtriva Prescribing Information, p.4. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
15. Gilead Sciences - Emtriva Prescribing Information, p.8. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
16. Gilead Sciences - Emtriva Prescribing Information, p.9. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
17. Gilead Sciences - Emtriva Prescribing Information, p.12. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
18. Gilead Sciences - Emtriva Prescribing Information, p.13. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
19. Gilead Sciences - Emtriva Prescribing Information, p.13. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
20. USP Dictionary of USAN & Intern. Drug Names - 2003; p. 311
21. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus>. Accessed 11/26/03.
22. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus>. Accessed 11/26/03.
23. Merck Index - 2001, 13th Edition; p 630.
24. Merck Index - 2001, 13th Edition; p. 630
25. Merck Index - 2001, 13th Edition; p. 630
26. Gilead Sciences - Emtriva Prescribing Information, p.1. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
27. Gilead Sciences - Emtriva Prescribing Information, p.1. Available at: <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.
28. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
29. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
30. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.

Emtricitabine



31. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
32. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
33. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
34. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
35. ChemIDplus. - Available at: <http://chem.sis.nlm.nih.gov/chemidplus/>. Accessed 11/26/03.
36. World Health Organization (WHO) - 1999; International Nonproprietary Names for Pharmaceutical Substances, p.9. Available at: http://www.who.int/druginformation/vol13num3_1999/listr42.pdf. Accessed 11/26/03.
37. Gilead Sciences - Emtriva Prescribing Information, p. 1. Available at <http://www.emtriva.com/fpi.pdf>. Accessed 07/03/03.